



DIABETES MANAGEMENT PLAN 2022

INSULIN PUMP




Australian Paediatric Society
 ISPAD school e- learning professional development
<https://www.t1d.org.au>

ISPAD Position Statement
 Type 1 Diabetes in Schools
www.ispad.org/news/news.asp?id=420540

This document outlines the consented medical management prescribed for this individual student with Type 1 Diabetes while under the care and supervision of school or pre-school personnel. These medical orders cannot be altered by a third party without parental consent.

Name of Student	
Date of Birth	
Parent 1 / carer name	
Parent 1 contact	
Parent 2 / carer name	
Parent 2 contact	
Diabetes Educator name	
Diabetes Educator contact	
Doctor name	
Doctor contact number	
Insulin Pump type / model	
CGM type /model	



The **Diabetes Action Plan** is a concise consented medical emergency response plan that outlines prescribed blood glucose target levels and urgent management of high and low blood glucose levels. A **Health Support Plan** should be developed between parent and school to clearly outline how the school will execute these medical orders.

Education and Training

Individualised training: Best practice is the development of a close working relationship between parent and school. The parent is best placed to understand the individual needs of their child so the parent should provide training of school personnel on those needs. The student’s diabetes team may also assist with training. With an increasing variety of diabetes technology available, individualised training is essential.

Education providers: are responsible to ensure that school personnel responsible for the health services of the student are performing duties within their scope of practice and with appropriate qualifications and training.

Parent/legal guardian: It is the parent’s right and responsibility to instruct the school on the specific care required for their child. The parent/legal guardian must be fully informed of any instruction, advice or training that is provided to school personnel regarding the needs of their child with Type 1 Diabetes.

The T1D diabetes e-learning courses for school staff (www.t1d.org.au) are the required professional development for this student’s individualised medical needs and must not be substituted by alternative learning courses.

- The school personnel with a direct supervision responsibility for this student (class teacher/ homeroom teacher/ special subject teachers) are requested to complete level 1 and level 2 T1D e-learning courses.
- The school personnel who are responsible for the fulfilment health services and administration of medication for this student must also complete T1D Level 3 e-learning course.

Parent..... Doctor.....Date /..... /2022

Daily requirements for school / preschool

Blood glucose (BG) check	Remind <input type="checkbox"/>	Observe <input type="checkbox"/>	Assist <input type="checkbox"/>	Perform <input type="checkbox"/>	No <input type="checkbox"/>
Blood glucose entry into pump	Remind <input type="checkbox"/>	Observe <input type="checkbox"/>	Assist <input type="checkbox"/>	Perform <input type="checkbox"/>	No <input type="checkbox"/>
Carbohydrate entry into pump	Remind <input type="checkbox"/>	Observe <input type="checkbox"/>	Assist <input type="checkbox"/>	Perform <input type="checkbox"/>	No <input type="checkbox"/>
Food Consumption	Remind <input type="checkbox"/>	Observe <input type="checkbox"/>			No <input type="checkbox"/>
Connect and disconnect pump line		Observe <input type="checkbox"/>	Assist <input type="checkbox"/>	Perform <input type="checkbox"/>	No <input type="checkbox"/>

Note - Insulin MUST always be administered before the student commences eating

Self-Management Skills

- Full Support:** All care performed by qualified/appropriately trained school staff
- Supervision:** Assistance and supervision by qualified /appropriately trained staff
- Self-care:** Student manages routine diabetes care requirements independently. Qualified/appropriately trained staff to provide support and supervision where required and as needed, in the event of becoming unwell.

The education provider's obligation to ensure the safety and wellbeing of the T1D student under their supervision is not diminished when the student may be capable of undertaking some routine diabetes care requirements. Qualified/appropriately trained school staff must be available to support and/or perform the health services required for the student.

Blood /Sensor Glucose checking times

Student must wash and dry hands (sanitiser gel/hand rub cause errors and are NOT recommended).
CGM readings may be substituted for blood glucose readings unless low or not consistent with symptoms **Yes/ No**

	Time	Blood (finger-prick)	Sensor (CGM/Flash)
Low glucose (Hypo) suspected	Any time	Yes / No	Yes / No
Exams	Prior to exam/during	Yes / No	Yes / No
Upon arrival to school		Yes / No	Yes / No
Pre-morning snack		Yes / No	Yes / No
Pre-morning recess		Yes / No	Yes / No
After morning recess		Yes / No	Yes / No
Pre-lunch		Yes / No	Yes / No
Pre-exercise / physical activity		Yes / No	Yes / No
After lunch recess		Yes / No	Yes / No
Pre-leaving school		Yes / No	Yes / No

Blood glucose levels vary with activity levels, stress, excitement, illness, menstruation, and food type/quantity.

Low blood glucose = less than _____mmol/l

Symptoms and the prescribed treatment are outlined in the Diabetes Action Plan.

Low glucose must be treated without delay with continuous responsible adult supervision during recovery.
Immediately notify parents for instruction if low glucose persists beyond 30 minutes despite treatment.

Low blood glucose ("Hypo") supplies are located:

In classroom under supervision **Yes / No** With student **Yes / No** Other _____

Please ensure low glucose kit is with the student when leaving school grounds on a school related activity.

Glucagon injection where prescribed is located (student locker is not recommended):

In classroom: **Yes /No** In "sick bay": **Yes / No** In Office: **Yes / No** Other _____

High blood glucose = greater than 8.0 mmol/l

The prescribed treatment is outlined in the Diabetes Action Plan.

Administer insulin "correction bolus" if this student's blood/sensor glucose is above _____ mmol/l or _____ mmol/l if physical activity is imminent or downward arrow on CGM. **Yes/No**

Pump will automatically administer insulin in response to high glucose levels. (Annexure 5) **Yes/ No**

Notify parents if the student is **unwell** with high blood glucose. **This may be life threatening.**

If in care of school for an extended time (e.g. overnight/ school camp) **blood ketones** must be tested if unwell. A level greater than 0.6mmol/l requires immediate medical management.

Physical Activity Strategy

Students with T1D should be encouraged and enabled to participate in physical activity with the appropriate adjustments for safety and optimal performance.

1. Insulin dose reduction before scheduled physical activity

Reduce insulin dose up to 2 hours before exercise **Yes / No**

This is achieved by reduced carbohydrate dosage by entering _____g less for food entries up to 2 hours before exercise

And /or by entering temporary basal rate of _____ for 2 hours before exercise and 1 hour after exercise.

2. Carbohydrate addition before and after exercise

Give _____ (without insulin) before exercise if Blood /Sensor Glucose is under 8 mmol/l or expected to be under 8 mmol/l within 15 minutes as indicated by downward trend arrow on CGM.

3. Swimming, vigorous activity and contact sports

An insulin pump should stay connected in most activities but may be disconnected from the student for up to 2 hours swimming, vigorous activity and contact sports. A disconnected pump **MUST** be handled with care by an adult and reconnected by the authorised adult or returned to the child at the end of activity.

Continuous Glucose Monitors (CGM)

Continuous Glucose Monitoring (CGM) is now a very common and useful part of the student's routine diabetes management. The CGM may read continuously or intermittently depending on the technology in use. (Annexure 4)

The following devices are used as usual part of the medical treatment and must always be present with the student during school hours for medical purposes and communication to parents:

Mobile Phone Yes / No **Smart Watch** Yes / No **Pump** Yes / No **Receiver** Yes / No

If the student is wearing any type of CGM, refer to Annexure 4. Please discuss the CGM plan and interventions with parents.

If the device reads low or the student has symptoms of low blood glucose, a finger prick blood glucose is required to confirm the result. **Symptoms of low blood glucose should be treated regardless.**

Sensor glucose (SG) devices have approximately a 5 to 15 minute lag time behind **blood glucose (BG)** levels.

Please use Trend Arrows on CGM (Annexure 4) to manage glucose levels **Yes / No**

Insulin Pump Troubleshooting Skills

Parental guidance must be sought for insulin pump and/or CGM issues relating to insulin delivery. Parents must be notified immediately if insulin pump delivery fails.

The parental assessment of their child's capabilities include that the student is:

- | | |
|---|----------|
| • Able to fill insulin reservoir and prepare tubing and cannula change | Yes / No |
| • Able to insert new cannula / infusion line | Yes / No |
| • Able to disconnect and reconnect tubing if required | Yes / No |
| • Able to self-administer insulin injection if required without supervision | Yes / No |
| • Able to test ketones and interpret results | Yes/ No |
| • Action and interpret CGM Alerts | Yes / No |

Coeliac Disease

This student also has coeliac disease so must avoid gluten (wheat) Yes / No

Record Keeping

All treatment / supervision of treatment undertaken must be recorded with the action taken, time and dose (where relevant) including (but not limited to):

- blood /sensor glucose results,
- insulin administration,
- incidence and treatment of low blood glucose ("hypos")

Parents may request this information to be recorded in a Communication Book or other daily advice document (Annexure 4). These documents are medical records and remain the property of the parent.

Predictive alert/suspend, Hybrid Closed Loop, Open Artificial Pancreas Systems

Technology innovations have created systems that automatically feed information from CGM to insulin pump, with the pump then automatically responding by adjusting insulin administration to the student.

This student uses the following system for diabetes management:

- Predictive Low Alert with Low Blood Glucose suspend
- Hybrid Closed Loop (Increases and decreases basal insulin)
- Advanced Hybrid Closed Loop (adjusts basal insulin and gives correction bolus based on serum glucose)
- Open Artificial Pancreas System

In these systems, diabetes management is different. The Diabetes Action Plans notates a modified response to low blood glucose levels. Details of the system being used by this student is provided in **Annexure 5**

The school is requested to assist the student to appropriately use this technology. The parent will supply the school with information and the required assistance to maintain optimal use of such technology while under the care of the school.

Why Target Glucose is Important

High blood glucose levels should NOT be accepted as commonplace and MUST be acted upon. High blood glucose levels at school are unacceptable. High blood glucose levels can cause:

- brain injury
- significant impact upon mood, concentration, learning, memory and verbal comprehension.
- markedly increased risk of long-term medical complications
- shortened life expectancy.

Communication

*Caring for a student with T1D is best achieved through a cooperative, supportive and respectful relationship between the three key stakeholders – **parent** (and student when they are capable of greater independence in self-care), **school personnel** and **medical team**.*

Parents are the final arbiters of whether their child can self-manage certain aspects of T1D, including glucose monitoring and self-administration of insulin. The medical team should guide and support parents to ensure the student is not subject to inappropriately unrealistic expectations.

An effective communication process between parent/student (when capable) and school personnel should be respectful, transparent and easily accessible.

Roles and Responsibilities

Medical Staff/ Treating Medical Team

The student's treating doctor or nurse practitioner is responsible for prescribing medications. The medical team is responsible for outlining in detail the recommended medical requirements for that student. This cannot be delegated to a third party that is not authorized or not suitably qualified.

School personnel should consider the student's medical team as an accessible resource to contact with parental permission. A single member of the medical team should be identified as the source of contact for each student.

Parent/Legal Guardians

Parents are ultimately responsible for the medical decisions made on behalf of their child. Therefore, the parent's informed consent and decisions regarding the health and well-being of their child are paramount. It is imperative that parents remain engaged as part of the team even when the student with T1D reaches adolescence.

The school must contact the parent/legal guardian in first instance, however if the parent is uncontactable or in the event of an emergency, school staff are authorised to contact our child's treating medical team.

The contact person from the student's medical team isPh.....

Privacy

This Diabetes Management Plan contains private and confidential medical orders and individual health information. This information cannot be shared with any 3rd party without specific written parental/ legal guardian consent.

Informed Consent

The contents of this Diabetes Management Plan are my prescribed treatment and individualised medical orders

Signed _____ (Doctor) Name _____ Date ____/____/2022

I agree with prescribed treatment as outlined in this Diabetes Management Plan, emergency response Action Plan and associated Annexures. I understand that

- I am responsible for supply of all Type 1 Diabetes information and material, equipment, insulin, hypoglycaemia supplies and Glucagon Hypo kit.
- I understand it is my right and responsibility to notify the school of any changes to the prescribed treatment and medical needs of my child with Type 1 Diabetes.
- I understand that the education provider/ school/ pre-school is responsible for the fulfillment of this medical order in accordance with their legal and regulatory obligations.

This plan and associated documents cannot be altered or amended without consent.

Signed _____ (parent / legal guardian) Date ____/____/2022

Name _____

Annexure 1

Terminology

- **Insulin Pump** is a pre-programmed computerised device that delivers insulin through an infusion set delivering constant background “basal” insulin and a calculated “bolus” insulin dose upon entry of blood/sensor glucose and /or carbohydrate food content.
- **Cannula** – small Teflon or metal tube that is inserted under the skin as the portal for insulin delivery. Must be replaced every 2-3 days. Teflon may kink causing failure to deliver insulin (line failure).
- **Reservoir / Cartridge** – plastic container within the pump containing enough insulin for 2-3 days.
- **Tubing** – plastic tubing connecting the cartridge to the cannula - it cannot kink.
- **Infusion set or “line”**– Cannula + cartridge + tubing.
- **Ketones** – chemicals produced by fat breakdown when glucose becomes unavailable as a fuel for cells to burn for energy (e.g. failure of insulin delivery). Small amounts of ketones are not usually a concern however when present in large amounts can induce nausea and vomiting, potentially leading to serious problems.
- **Temporary Basal** - an increase or decrease in insulin basal delivery for a prescribed length of time.
- **Extended/ dual/ combo wave** –extending an insulin bolus over an extended period of time.
- **Basal** – background insulin that is delivered continuously.
- **Bolus** – insulin administered prior to food to match carbohydrate content of food.
- **Correction Bolus**- insulin administered to correct a high blood glucose.
- **Line Failure** – disruption to the insulin administration and delivery which may be caused by cannula kinking, blocking or being dislodged. It is rarely caused by tubing issues.
- **Insulin Pump Delivery Failure** – failure of the pump to deliver and administer insulin as a result of a pump technical fault, pump switched off / suspended or Line Failure.

Vomiting and Type 1 Diabetes

Vomiting requires urgent assessment of blood glucose and blood ketones. It may indicate life-threatening “DKA”. Never assume the cause of vomiting in a person with Type 1 Diabetes to be “gastro”, food poisoning, migraine, excessive alcohol, until it is clear that insulin has been effectively administered. This will be evident if blood glucose is high and blood ketones are greater than 0.6mmol/l. Refer to **Diabetes Action Plan**.

ISPAD Recommended levels of school staff education and training

Level 1 - All school personnel should be educated about basic medical understanding of T1D (including recognition and urgency of treatment for low blood glucose) and the effect of T1D on the student and the entire family including the social, economic and emotional impact of living with T1D.

Level 2 - Those school personnel most responsible for the day-to-day management of the child with T1D should be also trained for **the individual student** to

1. recognize low blood glucose symptoms and signs,
2. initiate treatment for high or low blood glucose levels and
3. know and understand when and whom to call for assistance, including emergency responders, parents and medical team.

Level 3 -Those school personnel with authorisation or seeking authorisation through training and informed parental consent to administer insulin require a higher level of training on:

- insulin administration, including dose calculation and adjustments
- the legal aspects of insulin administration insulin
- delivery devices including insulin pumps
- glucagon administration

School camps

The medical management at school camp is no different to management on the school campus, with the same responsibilities, action plans, and target blood glucose levels. Special preparation including upskilling, risk assessment and communication strategies may be required to account for remoteness, activity, supplies and self-management skills. Detailed forward planning between school and parent (and sometimes the student’s medical team) is essential and will require a separate, individualised Health Support Plan for remote camps. The Diabetes Management Plan and Diabetes Action Plan MUST accompany the student on camp.

Annexure 2:

A Parent Guide – International best practice Type 1 Diabetes care in Australian schools

A Parent Guide, based on International Society for Pediatric and Adolescent Diabetes (ISPAD) guidelines, has been produced and endorsed by the Australian Paediatric Society to assist parents understand how they may access best practice Type 1 Diabetes management for their child at school. The clinical guidance is based on ISPAD standards and is consistent with the ISPAD principles of international best practice clinical governance, advocacy, education and science. Available to download at <https://www.t1d.org.au/diabetes-at-school/a-parent-guide>

School Personnel Training Agenda

An agenda to assist school personnel training by the parent has been developed by the Australian Paediatric Society. This is a checklist to assist coverage of all important diabetes topics during individualised training. Available to download at https://www.t1d.org.au/images/docs/T1DLC_Training_Agenda.pdf

General Issues with Type 1 Diabetes – ISPAD Position Statement

ISPAD, the world authority on contemporary diabetes management, states: *Schools are responsible for ensuring that their personnel are adequately educated about T1D and trained in the application of prescribed treatment for the individual student. The content of the training is the responsibility of the medical team and parent. Training should be executed by people with appropriate understanding of the student's individual needs and skill set.*

School personnel must understand the emotional burden experienced by families when given a diagnosis of an incurable disease such as T1D that will relentlessly impact upon the student, siblings, family relationships and parental working lives.

A diagnosis of T1D may cause students to feel different from peers and put them at risk of being stigmatised, resulting in a higher risk of experiencing anxiety and depression. The traumatised family may feel helpless and disempowered and yet have an obligation to advocate for their child.

Each family will have access to different resources, coping skills and economic circumstances. School personnel will have varying interest and levels of expertise. Hence care of the student must be individualized.

Schools should not expect that young people with diabetes will "learn responsibility" for self-managing T1D by leaving them unsupported during school hours. Nor will the duration the student has lived with T1D determine their ability to be self-sufficient. Young students may be capable but should not be solely responsible for their management at school. (ISPAD PS 6.9)

Young children are not capable of managing diabetes care. They **require extra support at school** and all very young children need full support to ensure safe and legal insulin delivery and other diabetes care. The child with diabetes may be encouraged to be involved in care and perform some tasks by themselves under supervision. The student may be capable but should not be responsible for Type 1 management during school hours as the effects of low or high blood glucose may seriously impair judgement.

There is no consensus as to what age the student may be expected to have responsibility for self-care during the school day. In most cases the child is mature enough by 12 years but a neurocognitive dysfunction, learning disability or psychosocial vulnerability can cause prolonged need for support. The parent is the best and most appropriate person to judge this in conjunction with the child's medical team and should document the amount of assistance and supervision required in the child's individual Diabetes Management Plan.

There is increasing recognition that adolescents are generally not capable of total diabetes care until they leave school and their forebrain fully develops. Adolescents have other interests, do not want to be different from their peers and having a condition such as diabetes may carry a stigma, so diabetes management is often not a high priority. Diabetes teams aim to encourage children with Type 1 to enjoy active "normal" lives not inhibited by Type 1. Discrimination, exclusion, inappropriate comments, and lack of facilitation of Type 1 requirements during school time for many children can destroy such ethos.

Annexure 4: Continuous Glucose Monitor (CGM)

There are two main systems of CGM in Australia:

1. **Real-time CGM** (rtCGM) utilize real-time alarms for thresholds and predictions of hypo- and hyperglycaemia, as well as rate of change alarms for rapid glycaemic excursion. Some CGM sensors to transmit signals to the “cloud,” and allow for digital remote monitoring, through which caregivers are able to view a patient's CGM tracing and receive alerts on their own devices, including smartphones, tablets, and smart watches.
2. **Intermittently scanned” CGM** (isCGM) systems, also known as flash glucose monitoring (FreeStyle Libre), do not automatically display glucose readings at regular intervals, but report glucose levels only when the user scans the sensor by holding a reader, or a mobile phone, close to the sensor. (5)

Each CGM available has different options and platforms to access and receive the sensor glucose data, notifications, alerts and alarms from the CGM transmitter via Bluetooth. These include, a specific data receiver, an application on a mobile (smart) phone, smart watches, web pages and direct to the insulin pump screen.

Some CGM technologies allow the data to be accessed from multiple platforms while others have a single specific platform to receive and access data. When the student’s CGM data is “shared” it is done by an application using a wireless network or cellular data. Hence some CGM technologies enable the student to share the real time glucose monitoring data with others, who might include the school nurse, authorised school personnel and the parent. The ability for others to view the glucose data and receive the notifications, alerts and alarms from the individual with T1D is referred to as “remote monitoring”.

CGM provides valuable information about glucose levels for the student, caregivers, school nurse, and diabetes care team. CGM update glucose data every 5 minutes, providing 288 readings per day. In addition, CGM have trend arrows, that in combination with the current glucose level, allow the student, and the school personnel responsible for the student’s complex medical care, to know what the current glucose level is, where it is going, and how fast it is changing.

Dexcom CGM	Libre	Medtronic CGM	Significance	Prevent low by (consider the effect of exercise)
			BG will fall >2.5mmol/l in 15 mins	If BG 6.5 mmol/l or lower – treat as low per Concise Action Plan
			BG will fall >1.7 mmol/l in 15 minutes	If BG 5.7 mmol/l or lower treat as low per Concise Action Plan
			BG will fall >0.8mmol/l in 15 minutes	If BG 4.8mmol/l or lower treat as low per Concise Action Plan
			BG will fall <0.8mmol/l in 15 minutes	Observe

Annexure 5: Insulin Pumps

Insulin Pump/ CGM interaction

There are an increasing number of students who manage diabetes with advanced technology that will be increasingly used in the school environment. These technologies are dependent upon CGM sensor glucose readings and include:

- Predictive Low Alerts with Low Glucose Suspend (the insulin pump will cease delivering background insulin if the sensor glucose predicts a low glucose levels).
- Hybrid Closed Loop - the pump will suspend insulin if predicting low sensor glucose and administer additional insulin if predicting high sensor glucose. Food (carbohydrates) must still be entered into the insulin pump prior to eating.
- Open Artificial Pancreas System -

The school is requested to assist the student to appropriately use this technology. The parent will supply the school with information about these systems and the required assistance to maintain optimal use of such technology during school hours.

Common Advanced Systems

Tandem T Slim

Basal IQ - Pairs with Dexcom G6 CGM to enable the pump to suspend basal insulin delivery if sensor glucose is predicted to fall low and resumes basal delivery automatically when the serum glucose rises.

Food (carbohydrates) must still be entered into the insulin pump prior to eating.

Because pump suspends basal insulin delivery, do not overtreat low blood glucose. See Action Plan.

Control IQ - Pairs with Dexcom G6 CGM and connects with support phones to share data. Automode adjusts basal insulin based on sensor glucose levels and will deliver a correction insulin bolus every hour as required.

Medtronic 670G and 770G

670G pairs with the Medtronic Guardian Link (3) CGM

770G pairs with the Medtronic Bluetooth Guardian Link (3) CGM and connects with support phones to send information via the student's phone data.

Auto Mode: Adjusts basal insulin continuously based on sensor glucose.

Must be calibrated to maintain hybrid closed loop status.

If alarms sound, follow the prompts on the screen.

Food (carbohydrates) must still be entered into the insulin pump prior to eating.

Because the pump suspends basal insulin delivery, do not overtreat hypoglycaemia- see action plan

Manual mode: When the pump is in manual mode, the **basal low glucose suspend must be turned on** and the pump will suspend basal insulin if the serum glucose is predicted to go low.

Medtronic 780G

Pairs with the Medtronic Bluetooth Guardian Link (3) continuous glucose monitor and connects to support phones via the child's phone data.

Auto Mode: the pump adjusts basal insulin (like 770G) and also gives a correction bolus of insulin every 5 minutes if the blood glucose is above 6.7 mmol/l and the basal is on maximum delivery.

Food (carbohydrate) must still be entered into the pump and the blood/sensor glucose entered with the food bolus.

When not adding food, the pump should automatically correct by delivering insulin based on the serum glucose readings in the CGM.

Hypoglycaemia -Because the pump suspends basal insulin delivery, do not overtreat hypoglycaemia- see action plan

Annexure 6: Other Individual Requirements

The following are also required for the complex care of my child with Type 1 Diabetes to maintain blood glucose levels as much as possible in the normal range whilst under the care of school:

Signed..... (parent)

Date / / 2022